

## List of Select Studies Supporting the Science Behind DietCypher

Goni L, Sun D, Heianza Y, et al. A circadian rhythm-related MTNR1B genetic variant modulates the effect of weight-loss diets on changes in adiposity and body composition: the POUNDS Lost trial. *Eur J Nutr.* 2019;58(4):1381-1389.

Hotta K, Nakata Y, Matsuo T, et al. Variations in the FTO gene are associated with severe obesity in the Japanese. *J Hum Genet.* 2008;53(6):546-553.

Kamura Y, Iwata M, Maeda S, et al. FTO Gene Polymorphism Is Associated with Type 2 Diabetes through Its Effect on Increasing the Maximum BMI in Japanese Men. *PLoS One.* 2016;11(11):e0165523.

Liu X, Zhang F, Zhang Y, et al. PPM1K Regulates Hematopoiesis and Leukemogenesis through CDC20-Mediated Ubiquitination of MEIS1 and p21. *Cell Rep.* 2018;23(5):1461-1475.

Loos RJ, Yeo GS. The bigger picture of FTO: the first GWAS-identified obesity gene. *Nat Rev Endocrinol.* 2014;10(1):51-61.

Mattei J, Qi Q, Hu FB, Sacks FM, Qi L. TCF7L2 genetic variants modulate the effect of dietary fat intake on changes in body composition during a weight-loss intervention. *Am J Clin Nutr.* 2012;96(5):1129-1136.

Merritt DC, Jamnik J, El-Sohemy A. FTO genotype, dietary protein intake, and body weight in a multiethnic population of young adults: a cross-sectional study. *Genes Nutr.* 2018;13:4.

Nagy K, Fiatal S, Sándor J, Ádány R. Distinct Penetrance of Obesity-Associated Susceptibility Alleles in the Hungarian General and Roma Populations. *Obes Facts.* 2017;10(5):444-457.

Qi Q, Bray GA, Smith SR, Hu FB, Sacks FM, Qi L. Insulin receptor substrate 1 gene variation modifies insulin resistance response to weight-loss diets in a 2-year randomized trial: the Preventing Overweight Using Novel Dietary Strategies (POUNDS LOST) trial. *Circulation.* 2011;124(5):563-571.

Roswall N, Ångquist L, Ahluwalia TS, et al. Association between Mediterranean and Nordic diet scores and changes in weight and waist circumference: influence of FTO and TCF7L2 loci. *Am J Clin Nutr.* 2014;100(4):1188-1197.

Rzehak P, Scherag A, Grallert H, et al. Associations between BMI and the FTO gene are age dependent: results from the GINI and LISA birth cohort studies up to age 6 years. *Obes Facts.* 2010;3(3):173-180.

Sacks FM, Bray GA, Carey VJ, et al. Comparison of weight-loss diets with different compositions of fat, protein, and carbohydrates. *N Engl J Med.* 2009;360(9):859-873.

Ślęzak R, Leszczyński P, Warzecha M, Łaczmański Ł, Misiak B. Assessment of the FTO gene polymorphisms in male patients with metabolic syndrome. *Adv Clin Exp Med*. 2018;27(11):1581-1585.

Sotos-Prieto M, Smith CE, Lai CQ, Tucker KL, Ordovas JM, Mattei J. Mediterranean Diet Adherence Modulates Anthropometric Measures by TCF7L2 Genotypes among Puerto Rican Adults. *J Nutr*. 2020;150(1):167-175.

Wang L, Shen H, Liu H, Guo G. Mixture SNPs effect on phenotype in genome-wide association studies. *BMC Genomics*. 2015 Feb 3;16(1):3.

Xu M, Qi Q, Liang J, et al. Genetic determinant for amino acid metabolites and changes in body weight and insulin resistance in response to weight-loss diets: the Preventing Overweight Using Novel Dietary Strategies (POUNDS LOST) trial. *Circulation*. 2013;127(12):1283-1289.

Zhang X, Qi Q, Zhang C, et al. FTO genotype and 2-year change in body composition and fat distribution in response to weight-loss diets: the POUNDS LOST Trial [published correction appears in *Diabetes*. 2013 Feb;62(2):662. Smith, Steven R [added]; Bray, George A [added]]. *Diabetes*. 2012;61(11):3005-3011.